



Available online at www.sciencedirect.com



Journal of Electromyography and Kinesiology xxx (2004) xxx-xxx

JOURNAL OF
ELECTROMYOGRAPHY
AND
KINESIOLOGY

www.elsevier.com/locate/jelekin

Quantification the relationship between FITA scores and EMG skill indexes in archery

H. Ertan ^{a,*}, A.R. Soylu ^b, F. Korkusuz ^a

^a Physical Education and Sports Department, Middle East Technical University, 06531, Turkey

^b Biophysics Department, Hacettepe University Medical Faculty, 06532, Turkey

Received 23 January 2004

Abstract

Forearm electromyographic (EMG) data are assumed to be an effective method in estimating performance level in archery. The aim of the current study was to establish archery skill indexes based on EMG data. Elite ($n = 7$, FITA score = 1303.4 ± 26.2), beginner ($n = 6$, FITA score = 1152 ± 9.0) and non-archers ($n = 10$, assumed FITA score = 250 0), were involved in the study. EMG activity of Muscle flexor digitorum superficialis and Muscle extensor digitorum were quantified. Two-second periods – 1 s before and 1 s after the fall of the clicker – were used to obtain averaged and rectified EMG data. The averaged and rectified EMG data were filtered by averaging finite impulse response filter with 80 ms time window and then normalized with respect to maximum voluntary contraction. To estimate FITA scores from EMG data, the following skill indexes that based on mean area under some parts of processed EMG waveforms was offered for archery. These were the pre-clicker archery skill index (PreCASI), post-clicker archery skill index (PostCASI), archery skill index (ASI) and post-clicker archery skill index 2 (PostCASI2). The correlations between rank of FITA scores and natural logarithms of archery skill indexes were significant for $\log(\text{PreCASI})$: $r = -0.66$, $p < 0.0008$; for $\log(\text{PostCASI})$: $r = -0.70$, $p < 0.0003$; for $\log(\text{ASI})$: $r = -0.74$, $p < 0.0001$; $\log(\text{PostCASI2})$: $r = -0.63$, $p < 0.002$. It is concluded that EMG skill indexes may be useful for: (a) assessing shooting techniques, (b) evaluation of archers' progress and (c) selection of talented archers.

© 2004 Elsevier Ltd. All rights reserved.

Keywords: Archery; Electromyography; Muscular analysis and archery skill indexes

1. Introduction

Skill in archery is defined as the ability to shoot an arrow to a given target in a certain time span with accuracy [10]. Archery shooting is described as a three-phase movement as drawing, aiming and release [14]. Each of these phases represents a stable sequence of the collective movements and is ideal for studying the motor control and skill-acquired. An archer pushes the bow with an extended arm, which is statically held in

the direction of the target, where the other arm exerts a dynamic pulling of the bowstring from the beginning of the drawing phase, until the release is dynamically executed [10]. The release phase must be well balanced and highly reproducible to achieve commendable results in an archery competition [14].

The bowstring is released when audible impetus is received from a device called "clicker". Each arrow can be drawn to the exact distance and a release can be obtained and maintained by this device. The clicker is reputed to improve the archer's score and is used by all target archers [10]. The archer should react to the clicker as quickly as possible and synchronize the muscle activity of the whole body to attain eventual opti-

* Corresponding author. Tel.: +90 3122 104025; fax: +90 3122 101255.

E-mail address: hertan@metu.edu.tr (H. Ertan).

mal accuracy. In particular, there should be a repeated contraction and relaxation in the forearm and pull finger muscles during archery training and competitions according to the high number of arrows. When the clicker signal is heard, the archer relaxes the flexor group muscles of the forearm and actively contracts the extensor group muscles for producing the release. The archer should produce a muscular coordination between the forearm agonist and antagonist muscles [2,4,6,14].

Forearm muscles have special importance in holding and drawing the string by three-finger hook, taking it to full draw and releasing it. Importance of the activation patterns of the forearm muscles has been clarified and it is suggested that the archers, in different performance level, have different forearm contraction strategies. Elite and beginner archers displayed a stable contraction pattern before the clicker's impetus, while the non-archers showed a slight rising slope of contraction indicating an increase in the activity of the specified muscle groups before reaching the peak level [4].

Muscle co-activation is the simultaneous activity of various muscles acting around the joint. Evaluating the impact of skill level on the muscle activation/co-activation patterns is not a new topic [1,7,11]. The latency of antagonist EMG burst was strongly correlated with parameters of the first agonist EMG burst. For the movements described by the speed-insensitivity strategy, the quantity of both antagonist and agonist muscle activity can be uniformly associated with selected kinetic measures that incorporate muscle force–velocity relations. For movements collectively described by the speed-sensitivity strategy, no single rule can describe all the combinations of agonist–antagonist coordination that are used to perform these diverse tasks [5]. However, the question of whether an archer's performance level can be estimated by analyzing forearm EMG data is still unanswered. So, the aim of the current study was to quantify the relationship between some skill indexes and EMG measures by using linear regression.

2. Methods

2.1. Subjects

Three groups, (i) elite ($n = 7$, FITA score = 1303.4 ± 26.2), (ii) beginners ($n = 6$, FITA score = 1152 ± 9.0) and (iii) non-archers ($n = 10$, assumed FITA score = 250 ± 0), were involved in the study. The FITA score is a summation of four distances (for female: 70, 60, 50 and 30 m; for male: 90, 70, 50 and 30 m), which are set by the International Archery Federation (FITA). An archer shoots 36 arrows to each distance. So, he/she shoots totally 144 arrows in a FITA round where the

highest score can be 1440. The first group consisted of national team archers. Beginner archers from the city archery club formed the second group. The third group included university students with no background knowledge or experience on archery. The first and second groups had their FITA scores from official competitions. However, the non-archers group had not taken part any archery competition. To avoid the risks and dangers of non-archers' shoot, we did not have measured FITA scores for non-archers group. An interview was applied to four archery experts, who were working as national team coaches, for searching information about the highest score that a person who did not have any archery experience can shoot. They assigned 250 mean FITA score for a non-archer. Besides, the authors of this paper involved the students of a beginning archery class who had 8 weeks learning sessions to shoot a FITA round. These subjects had shot a mean of 288 FITA score (standard deviation 65). According to the findings of the interviews and the students' scores, the assignment of 250 was chosen as an appropriate FITA score for a beginner.

2.2. Electromyographic recordings

The measurement sites were prepared first by shaving the area and then lightly abrading and cleansing the skin with alcohol. Skin tack F55 circular Ag/AgCl surface electrodes, filled with conductive electrolyte, were then positioned longitudinally along each muscle. The center-to-center distance between two electrodes was approximately 2 cm. The reference electrode was placed on the olecranon process of the ulna of the drawing arm. The signals were pre-amplified (analog differential amplifiers, preamplifier gain 500), filtered using a bandpass filter (8–500 Hz), sampled at 1000 Hz, and converted in digital form by a 12 bit A/D converter.

Each subject participated in a single test session. EMG activity of the M. flexor digitorum superficialis (MFDS) and the M. extensor digitorum (MED) were quantified. Since MFDS and MED were surrounded by the other muscles closely, cross-talk effect may have been occurred during EMG measurements. It should be noted that the activity of flexors and extensors of the forearm measured as a whole in the current study. However, the EMG activities recorded from forearm muscles was named as activities of MFDS and MED for the sake of the specification of the mentioned finger movement pattern and referring to the recording sites. The surface electrodes were placed on the central portion of the each muscle. Palpating the selective muscles while subjects simulated the preparatory shooting position and performed maximum isometric contraction of these muscles recording sites on the drawing arm were identified.

155 2.3. Procedure

156 Forearm muscular activation strategies were evalu-
 157 ated from the EMG recordings immediately before
 158 and after the clicker's impetus. The arrow was initially
 159 positioned between the unattached end of the clicker
 160 and the bow-grip. As the arrow was pulled beyond the
 161 clicker, the clicker-lever fell on the bow-handle, which
 162 conveyed the signal to the archer that the arrow was
 163 appropriately positioned and is ready to be released.

164 A mechanical switch was attached under the clicker
 165 to accurately measure the point of the audible impetus.
 166 This audible impetus was superimposed with the EMG
 167 results in the same time frame. EMG recordings were
 168 for 5 s; 2.5 s prior and 2.5 s after the clicker's audible
 169 impetus. This time period included the last seconds of
 170 Full Draw, Aiming and the first seconds of Release
 171 and Follow Through phases. Two-second periods – 1 s
 172 before and 1 s after the fall of the clicker – were used
 173 to obtain averaged and rectified EMG data. The aver-
 174 aged [12] and rectified EMG data were filtered (averag-
 175 ing finite impulse response filter with 80 ms time
 176 window) and then normalized according to Clarys et
 177 al. [2].

178 Prior to the shootings, the maximum voluntary con-
 179 traction (MVC) of the MED and MFDS of each subject
 180 were determined on the basis of EMGs. Subjects con-
 181 tracted these muscles to the highest level against a stable
 182 resistance by forming three-finger hook as they did in
 183 holding the bowstring and MVC was obtained under

184 these circumstances. The angle between the proximal
 185 and distal interphalangeal joint was not changed during
 186 the isometric contractions of the mentioned muscle
 187 groups. EMG amplitudes were normalized with respect
 188 to MVC, i.e., they were expressed as percentages of
 189 MVC. By MVC normalization method, variations in
 190 the relationship could be found in the same muscle
 191 among subjects [3]. Figs. 1 and 2 show processed (aver-
 192 aged, rectified, filtered and normalized) EMG data for
 193 each group separately.

194 Each subject completed three trial shots to acquaint
 195 with the measurement conditions. Muscle activity was
 196 sampled for a 5-s period as the subjects completed 12
 197 successive shots. After elimination of noisy ones, aver-
 198 age of 8–12 single shots for each subject was used for
 199 calculations. For the shooting trials, the sampling was
 200 manually triggered shortly after the archer achieved a
 201 full (optimal) draw position, so that the release of the ar-
 202 row occurred at approximately the midpoint of the sam-
 203 pling period.

204 2.4. Processing of skill indexes

205 To estimate FITA scores from EMG data, the follow-
 206 ing skill indexes was offered for archery: pre-clicker ar-
 207 chery skill index (PreCASI), post-clicker archery skill
 208 index (PostCASI), archery skill index (ASI) and post-
 209 clicker archery skill index 2 (PostCASI2). For
 210 calculation of archery skill indexes, mean areas (area cor-
 211 responding to time interval/the time interval) under the




Fig. 1. Filtered and normalized EMG data (%MVC vs. ms): (a) average of elite archers, (b) average of beginner archers, (c) average of non-archers. Bold and thin lines correspond to MED and MFDS respectively. Clicker falls at zero.




Fig. 2. The EMG data in Fig. 1 are regroupped (MVC% vs. ms): (a) MED, (b) MFDS. Bold, medium and thin lines correspond to elite, beginner and non-archers, respectively. Clicker falls at zero.

212 processed EMG data of MFDS and MED were used.
 213 Assuming clicker falls at 0th ms, mean areas under pro-
 214 cessed EMG data of MED were calculated: 300 ms contin-
 215 uous time interval (from -220 to +80 ms), 280 ms
 216 continuous time interval (from +160 to +440 ms) and
 217 580 ms time interval (from -220 to +80 ms and from
 218 +160 to +440 ms) were accepted as PreCASI, PostCASI
 219 and ASI respectively. Mean area, 200 ms continuous time
 220 interval (from +240 to +440 ms), under processed EMG
 221 data of MFDS was also calculated as PostCASI2. Selec-
 222 tion of time intervals was based on maximisation of
 223 square of correlation coefficients (r^2) between the rank
 224 of the FITA scores and archery skill indexes. This maxi-
 225 misation gave four different archery skill indexes' time
 226 intervals and the relationship between rank of FITA
 227 scores and natural logarithms of skill indexes (log(Pre-
 228 CASI), log(PostCASI), log(ASI) and log(PostCASI2))
 229 were calculated using regression analysis (Fig. 3).

230 3. Results

231 Results of Kolmogorov-Smirnov test showed non-
 232 normal distribution for FITA scores, so rank of FITA
 233 scores were used in regression analysis by assigning rank
 234 numbers from 1 to 23 to every member of ascending or-
 235 dered FITA scores. If there were ties, rank of equal
 236 FITA scores was replaced by mean rank values. Since
 237 the relations between rank of FITA scores and archery
 238 skill indexes were found exponential, natural logarithms
 239 of indexes were used for making the relations linear and
 240 increasing estimation accuracy.

The correlations between rank of FITA scores and log
 of archery skill indexes were significant for log(PreCASI):
 $r = -0.66$, y -intercept = -5.43 , gradient = -6.05 , $n = 23$,
 $p < 0.0008$; for log(PostCASI): $r = -0.70$, y -intercept =
 -3.40 , gradient = -4.96 , $n = 23$, $p < 0.0003$; for log(A-
 SI): $r = -0.74$, y -intercept = -7.70 , gradient = -6.72 ,
 $n = 23$, $p < 0.0001$; log(PostCASI2): $r = -0.63$, y -inter-
 cept = -9.57 , gradient = -4.38 , $n = 23$, $p < 0.002$. See
 Fig. 3 for regression lines.

4. Discussion

Archery shooting is described as a three-phase (draw-
 ing, aiming and release) movement. Moreover, Nishiz-
 ono et al. [14] further divides the stages of a shot into
 six: bow hold, drawing, full draw, aiming, release and fol-
 low-through. PreCASI (300 ms continuous time interval
 from -220 to +80 ms) corresponds with the full draw,
 aiming and release movements. PostCASI (280 ms con-
 tinuous time interval from +160 to +440 ms) includes
 the initiation of the response to fall of the clicker. So,
 the second index covers the release, which is the begin-
 ning of the response to a given stimulus (fall of the click-
 er), and the follow-through, which is the completion of
 the specific movement pattern or muscle contraction,
 phases. 200-ms continuous time interval from +240 to
 +440 ms under processed EMG data of MFDS was also
 calculated as PostCASI2. It is mainly related with the
 completion of muscular activity or the changes in the
 position in metacarpophalangeal, proximal and distal
 interphalangeal joints. ASI (580 ms time interval: from

241
242
243
244
245
246
247
248
249

250

251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269




Fig. 3. Relationship between rank of FITA scores and natural logarithms of PreCASI (a), PostCASI (b), ASI (c) and PostCASI2 (d). Regression parameters and lines are shown on the graphs.

270 -220 to +80 ms and from +160 to +440 ms) is the
 271 weighted sum of PreCASI and PostCASI. It includes
 272 the fore period before the fall of the clicker, premotor
 273 and motor times, which is the time gap that the impulse
 274 is transmitted from the sensory organs to the central
 275 nervous system and then to the muscles [9]. The interval
 276 from the stimulus to the first muscular contraction re-
 277 corded in EMG is termed premotor Reaction Time and

is thought to represent the central nervous system pro- 278
 cesses. This index also covers the interval from the first 279
 change in EMG to finger movement, which is termed mo- 280
 tor Reaction Time and represents the processes associ- 281
 ated with the muscle itself [8,9,15-17]. The interval 282
 from the initiation of the response (the end of Reaction 283
 Time) to the completion of the movement is also involved 284
 in ASI. The archery indexes summarize the whole archery 285
 shooting movement. They separate the muscular activa- 286
 tion into small but meaningful pieces. They provide 287
 information on the whole muscular activation patterns 288
 before and after the fall of the clicker. 289

Archery shooting has a stable sequence and it includes 290
 the stance, holding, drawing, full drawing, aiming, releas- 291
 ing and follow-through movements. Each of these phases 292
 represents a stable sequence of the collective move- 293
 ments. An archer is supposed to hold the string by 294
 three-finger hook in the drawing hand [13]. Forming 295
 three-finger hook is an example of isometric contraction. 296
 So, the archer should not change the angles of the draw- 297
 ing fingers or proximal and distal interphalangeal joint 298
 until the release. Increasing or decreasing areas under 299
 the processed EMG data may be an indication of the 300
 changes in phalangeal joint angles before the fall of the 301
 clicker. That can be the reason why the rank of FITA 302
 scores has negative correlation with EMG activities. 303
 The archer tries to hold and carry the weight of the string 304
 by contracting the forearm muscles. On the contrary, a 305
 decrease in EMG activities in the forearm muscles indi- 306
 cates that the archer may disperse the weight of the string 307
 on the forearm, arm, shoulder girdle, and some of the 308
 back muscles. So, the percentage of the forearm muscles 309
 decreases in the collective archery shooting movement. 310

Having negative significant correlation between rank 311
 of FITA scores and log of archery skill indexes shows 312
 that increase in archery experience causes a decrease in 313
 area under the processed EMG data. The amplitude of 314
 the peak (Fig. 2(a), amplitude of peak around 260 ms) 315
 after the fall of the clicker is also decreases in line with 316
 the increase in archery experience. 317

The proposed archery skill indexes summarize the arch- 318
 ery shooting movement from beginning to the end. 319
 They involve and describe all of the shooting phases 320
 from the drawing hand aspect. Moreover, they expose 321
 a negative relationship between the forearm muscular 322
 activation and FITA scores. EMG skill indexes may 323
 be helpful evidences during the construction and trying 324
 out of new shooting techniques in archery in such a 325
 way that revision of the shooting technique can be based 326
 on this evidence. They seek specific information about 327
 the shooting technique as well as judgements or opinions 328
 of a coach, who can make only visual inspection of the 329
 technique. Those archery indexes and judgements by the 330
 archery coach together may serve as feedback for FITA 331
 score or in general performance improvement. EMG 332
 skill indexes may be useful for assessing shooting 333

334 techniques while they are being developed, to help shape
 335 them in their final forms. This may be applicable to all
 336 sorts of archery performance level (e.g., beginning or
 337 elementary level; national level; or world-class archer).
 338 It is concluded that EMG skill indexes may be useful
 339 for: (a) assessing shooting techniques, (b) evaluation of
 340 archers' progress and (c) selection of talented archers.

341 Acknowledgements

342 Middle East Technical University Scientific Research
 343 Grant number BAP 2002-050403 funded this study. The
 344 authors thank to the president and secretary general
 345 Prof. Dr. Uğur Erdener and Elif Ekşi of the Turkish Ar-
 346 chery Federation, respectively.

347 References

- 348 [1] A. Buccolieri, L. Avanzino, C. Trompetto, G. Abbruzzese,
 349 Relaxation in distal and proximal arm muscles: a reaction time
 350 study, *Clinical Neurophysiology* 114 (2) (2003) 313–318.
 351 [2] J.P. Clarys, J. Cabri, E. Bollens, R. Sleenckx, J. Taeymans, M.
 352 Vermeiren, R.G. Van, G. Voss, Muscular activity of different
 353 shooting distances, different release techniques, and different
 354 performance levels, with and without stabilizers, in target archery,
 355 *Journal of Sport Sciences* 8 (1990) 235–257.
 356 [3] C.J. De Luca, The use of surface electromyography in biome-
 357 chanics, *Journal of Applied Biomechanics* 13 (2) (1997) 135–163.
 358 [4] H. Ertan, B. Kentel, S.T. Tümer, F. Korkusuz, Activation
 359 patterns in forearm muscles during archery shooting, *Human*
 360 *Movement Science* 22 (2003) 37–45.
 361 [5] G.L. Gottlieb, M.L. Latash, D.M. Corcos, T.J. Liubinskas, G.C.
 362 Agarwal, Organizing principles for single joint movements: V.
 363 Agonist-antagonist interactions, *Journal of Neurophysiology* 67
 364 (6) (1992) 1417–1427.
 365 [6] M.P. Hennessy, A.W. Parker, Electromyography of arrow release
 366 in archery, *Electromyography and Clinical Neurophysiology* 30
 367 (1990) 7–17.
 368 [7] E. Kellis, F. Arabatzi, C. Papadopoulos, Muscle co-activation
 369 around the knee in drop jumping using the co-contraction index,
 370 *Journal of Electromyography and Kinesiology* 13 (3) (2003) 229–
 371 238.
 372 [8] R. Kerr, *Psychomotor Learning*, Philadelphia Saunders College
 373 Publication, 1982.
 374 [9] M.L. Latash, *Neurophysiological Basis of Movement*, Human
 375 Kinetics Pub., Champaign, IL, 1998.
 376 [10] P. Leroyer, V. Hoecke, N. Helal, Biomechanical study of the final
 377 push-pull in archery, *Journal of Sport Sciences* 11 (1993) 63–69.
 378 [11] V. Linnamo, T. Moritani, C. Nicol, P.V. Komi, Motor unit
 379 activation patterns during isometric, concentric and eccentric
 380 actions at different force levels, *Journal of Electromyography*
 381 *Kinesiology* 13 (1) (2003) 93–101.

- [12] P.E. Martin, W.L. Siler, D. Hoffman, Electromyographic analysis
 382 of bow string release in highly skilled archers, *Journal of Sport*
 383 *Sciences* 8 (1990) 215–221.
 384 [13] W. McKinney, M. McKinney, *Archery*, eighth ed., Brown &
 385 Benchmark, Madison, WI, 1997.
 386 [14] A. Nishizono, H. Shibayama, T. Izuta, K. Saito. Analysis of
 387 archery shooting techniques by means of electromyography, in:
 388 International Society of Biomechanics in Sports. Proceedings.
 389 Symposium V, Athens, Greece, 1987.
 390 [15] J.B. Oxendine, *Psychology of Motor Learning*, Appleton-Century-
 391 Crofts, New York, 1968.
 392 [16] R.A. Schmidt, *Motor Learning and Performance*, Human Kinet-
 393 ics Books, Champaign, IL, 1991.
 394 [17] A.J. Vander, J.H. Sherman, D.S. Luciano, *Human Physiology*,
 395 fifth ed., McGraw-Hill Publication Company, 1990.
 396
 397



Hayri Ertan has received his B.S. from Hacettepe University School of Sports Sciences and Technology in 1997 and M.Sc from Middle East Technical University, Department of Physical Education and Sports. He is working on his PhD dissertation on archery. He has been making researches on electrophysiological measurement methods especially surface electromyography, auditory event-related potentials. He is also national team coach for Physically Disabled archers.



Abdullah Ruhi Soylu was born in 1964. He received his B.S. (Electrical and Electronics Engineering, 1993) from Bilkent University. He received his M.D. (1994) and Ph.D. (Biophysics, 2003) in Hacettepe University Medical Faculty (HUMF). He is currently a lecturer in HUMF, Biophysics Department. His research focuses on surface electromyography, signal processing and event-related potentials.



Feza Korkusuz his B.S. from Ankara University, Medical Faculty in 1986 and his M.D. from Gazi University on Orthopedic Surgery and Traumatology. He has been working as the head of the Department of Physical Education and Sports and Health and Caring Center at Middle East Technical University since 1997. He has researches on biomaterials, medicine and sports, sports physiology and traumatology.